arXiv:0803.3353v1 [math.RA] 24 Mar 2008

ON STRONGLY $g(x)$-CLEAN RINGS

LINGLING FAN † AND XIANDE YANG‡

†Department of Mathematics and Statistics, Memorial University of Newfoundland, St.John's, A1C 5S7, Canada

Email address: lingling@math.mun.ca

‡Department of Mathematics, Harbin Institute of Technology, Harbin, 150001, China

Email address: xiande.yang@gmail.com

1

Abstract. Let R be an associative ring with identity, $C(R)$ denote the center of R, and $g(x)$ be a polynomial in the polynomial ring $C(R)[x]$. R is called strongly $g(x)$-clean if every element $r \in R$ can be written as $r = s + u$ with $g(s) = 0$, u a unit of R, and $su = us$. The relation between strongly $g(x)$-clean rings and strongly clean rings is determined, some general properties of strongly $g(x)$-clean rings are given, and strongly $g(x)$-clean rings generated by units are discussed.

Key Words: strongly g(x)-clean rings, strongly clean rings, rings generated by units
Mathematics Subject Classification: 16U60, 16U99

1. Introduction

Let R be an associative ring with the group of units $U(R)$. R is called *clean* if for every element $r \in R$, $r = e + u$ with $e^2 = e \in R$ and $u \in U(R)$ [10] and R is called *strongly clean* if in addition, $eu = ue$ [11].

Let $C(R)$ denote the center of a ring R and $g(x)$ be a polynomial in $C(R)[x]$. Camillo and Simón [2] say R is $g(x)$-*clean* if for every element $r \in R$, $r = s + u$ with $g(s) = 0$ and $u \in U(R)$. If V is a countable dimensional vector space over a division ring D, Camillo and Simón proved that $End(_DV)$ is $g(x)$-clean if $g(x)$ has two distinct roots in $C(D)$ [2]. Nicholson and Zhou generalized Camillo and Simón's result by proving that $End(_RM)$ is $g(x)$-clean if $_RM$ is a semisimple R-module and $g(x) \in (x-a)(x-b)C(R)[x]$ where $a, b \in C(R)$ and $b, b-a \in U(R)$ [12]. [5, 14] completely determined the relation between clean rings and $g(x)$-clean rings independently. What is the relation between strongly clean rings and $g(x)$-clean rings?

Date: March 24, 2008.

[1‡] : Corresponding author

In this paper, we continue this topic. In Section 2, we define strongly $g(x)$-clean rings and determine the relation between strongly $g(x)$-clean rings and strongly clean rings; in Section 3, some general properties of strongly $g(x)$-clean rings are given; and in Section 4, some classes of strongly $g(x)$-clean rings generated by units are discussed.

Throughout the paper, $\mathbb{T}_n(R)$ denotes the upper triangular matrix ring of order n over R, $\mathbb{N}$ denotes the set of all positive integers, and $\mathbb{Z}$ represents the ring of integers.

2. STRONGLY $g(x)$-CLEAN RINGS VS STRONGLY CLEAN RINGS

Definition 2.1. Let $g(x) \in C(R)[x]$ be a fixed polynomial. An element $r \in R$ is *strongly $g(x)$-clean* if $r = s + u$ with $g(s) = 0$, $u \in U(R)$, and $su = us$. R is *strongly $g(x)$-clean* if every element of R is strongly $g(x)$-clean.

Strongly clean rings are exactly strongly $(x^2 - x)$-clean rings. However, there are strongly $g(x)$-clean rings which are not strongly clean and vice versa:

Let $\mathbb{Z}_{(p)} = \{\frac{m}{n} \in \mathbb{Q} : \gcd(p, n) = 1 \text{ and } p \text{ prime } \}$ be the localization of $\mathbb{Z}$ at the prime ideal $p\mathbb{Z}$ and C_3 be the cyclic group of order 3.

Example 2.2. *Let R be a commutative local or commutative semiperfect ring with $2 \in U(R)$. By the proof of* [14, Theorem 2.7]*, RC_3 is strongly $(x^6 - 1)$-clean. In particular, $\mathbb{Z}_{(7)}C_3$ is a strongly $(x^6 - 1)$-clean ring. Furthermore, by* [5, Example 2.2]*, $\mathbb{Z}_{(7)}C_3$ is strongly $(x^4 - x)$-clean. However, $\mathbb{Z}_{(7)}C_3$ is not strongly clean* [7, Example 1].

Example 2.3. *Let $R = \mathbb{Z}_{(p)}$ and $g(x) = (x - a)(x^2 + 1) \in C(R)[x]$. Then R is strongly clean but by a easy verification we know R is not strongly $g(x)$-clean. Let R be a boolean ring with more than two elements with $c \neq 0, 1$. Then R is strongly clean but R is not strongly $g(x) = (x + 1)(x + c)$-clean by* [5, Example 2.3].

However, for some type of polynomials, strong cleanness and strong $g(x)$-cleanness are equivalent.

Theorem 2.4. *Let R be a ring and $g(x) \in (x-a)(x-b)C(R)[x]$ with $a, b \in C(R)$. Then the following hold:*

1. *R is strongly $(x - a)(x - b)$-clean if and only if R is strongly clean and $(b - a) \in U(R)$.*

(2) *If R is strongly clean and $(b-a) \in U(R)$, then R is strongly $g(x)$-clean.*

Proof. (1). "$\Leftarrow$". Let $r \in R$. Since R is strongly clean and $(b-a) \in U(R)$, $\frac{r-a}{b-a} = e + u$ where $e^2 = e \in R$, $u \in U(R)$, and $eu = ue$. Thus, $r = [e(b-a)+a] + u(b-a)$ where $u(b-a) \in U(R), [e(b-a)+a-a][e(b-a)+a-b] = 0$, and $[e(b-a)+a]u(b-a) = u(b-a)[e(b-a)+a]$. Hence, R is strongly $(x-a)(x-b)$-clean.

"$\Rightarrow$". Since a is strongly $(x-a)(x-b)$-clean, there exist $u \in U(R)$ and $s \in R$ such that $a = s + u$ with $(s-a)(s-b) = 0$ and $su = us$. Hence, $s = b$. So $(b-a) \in U(R)$. Let $r \in R$. Since R is strongly $(x-a)(x-b)$-clean, $r(b-a)+a = s+u$ where $(s-a)(s-b) = 0$, $u \in U(R)$, and $su = us$. Thus, $r = \frac{s-a}{b-a} + \frac{u}{b-a}$ where $\frac{u}{b-a} \in U(R)$, $(\frac{s-a}{b-a})^2 = \frac{(s-a)(s-b+b-a)}{(b-a)^2} = \frac{(s-a)(b-a)}{(b-a)^2} = \frac{s-a}{b-a}$, and $\frac{s-a}{b-a} \cdot \frac{u}{b-a} = \frac{u}{b-a} \cdot \frac{s-a}{b-a}$. So R is strongly clean.

(2). By (1). □

Corollary 2.5. *For a ring R, R is strongly clean if and only if R is strongly (x^2+x)-clean.*

Proof. It follows from Theorem 2.4 by letting $a = 0$ and $b = -1$. □

Remark 2.6. The equivalence of strong (x^2+x)-cleanness and strong cleanness is a ring property since it holds for a ring R but it may fail for a single element. For example, $1 + 1 = 2 \in \mathbb{Z}$ is strongly clean but 2 is not strongly (x^2+x)-clean in $\mathbb{Z}$.

Example 2.7. *Let $C(X)$ denote the ring of all real-valued continuous functions from a topological space X to the real number field $\mathbb{R}$ and $C^*(X)$ denote the subring of $C(X)$ consisting of all bounded functions in $C(X)$. If X is strongly zero-dimensional, then $C(X)$ and $C^*(X)$ are strongly (x^2-nx)-clean rings for any $n \in \mathbb{N}$ since $C(X)$ and $C^*(X)$ are strongly clean* [1, 9] *and n is invertible in $C(X)$ and $C^*(X)$. If X is a P-space, then $\mathbb{M}_k(C(X))$ is strongly (x^2-nx)-clean for any $n, k \in \mathbb{N}$ because $\mathbb{M}_k(C(X))$ is strongly clean by* [6].

3. GENERAL PROPERTIES OF STRONGLY $g(x)$-CLEAN RINGS

Let R and S be rings and $\theta : C(R) \to C(S)$ be a ring homomorphism with $\theta(1) = 1$. For $g(x) = \Sigma a_i x^i \in C(R)[x]$, let $\theta^{'}(g(x)) := \Sigma\theta(a_i)x^i \in C(S)[x]$. Then θ induces a map $\theta^{'}$ from $C(R)[x]$ to $C(S)[x]$. If $g(x)$ is a polynomial with coefficients in $\mathbb{Z}$, then $\theta^{'}(g(x)) = g(x)$.

Proposition 3.1. *Let $\theta : R \to S$ be a ring epimorphism. If R is strongly $g(x)$-clean, then S is strongly $\theta^{'}(g(x))$-clean.*

Proof. Let $g(x) = a_0 + a_1x + \cdots + a_nx^n \in C(R)[x]$. Then $\theta^{'}(g(x)) = \theta(a_0) + \theta(a_1)x + \cdots + \theta(a_n)x^n \in C(S)[x]$. For any $s \in S$, there exists $r \in R$ such that $\theta(r) = s$. Since R is strongly $g(x)$-clean, there exist $t \in R$ and $u \in U(R)$ such that $r = t + u$ with $g(t) = 0$ and $tu = ut$. Then $s = \theta(r) = \theta(t) + \theta(u)$ with $\theta(u) \in U(S)$, $\theta^{'}(g(x))|_{x=\theta(t)} = 0$, and $\theta(t)\theta(u) = \theta(u)\theta(t)$. So S is strongly $\theta^{'}(g(x))$-clean. ☐

Corollary 3.2. *If R is $g(x)$-clean, then for any ideal I of R, R/I is $\overline{g}(x)$-clean with $\overline{g}(x) \in C(R/I)[x]$.*

Corollary 3.3. *Let R be a ring, $g(x) \in C(R)[x]$, and $1 < n \in \mathbb{N}$. If $\mathbb{T}_n(R)$ is strongly $g(x)$-clean, then R is strongly $g(x)$-clean.*

Proof. Let $A = (a_{ij}) \in \mathbb{T}_n(R)$ with $a_{ij} = 0$ and $1 \leq j < i \leq n$. Note that $\theta : \mathbb{T}_n(R) \to R$ with $\theta(A) = a_{11}$ is a ring epimorphism. ☐

Corollary 3.4. *Let R be a ring and $g(x) \in C(R)[x]$. If the formal power series ring $R[[t]]$ is strongly $g(x)$-clean, then R is strongly $g(x)$-clean.*

Proof. This is because $\theta : R[[t]] \to R$ with $\theta(f) = a_0$ is a ring epimorphism where $f = \sum_{i\geq 0} a_it^i \in R[[t]]$. ☐

Proposition 3.5. *Let $g(x) \in \mathbb{Z}[x]$ and $\{R_i\}_{i\in I}$ be a family of rings. Then $\prod_{i\in I} R_i$ is strongly $g(x)$-clean if and only if R_i is strongly $g(x)$-clean for each $i \in I$.*

Proof. It is clear by the definition and Proposition 3.1. ☐

For strongly clean rings, the author in [3, 4, 13] proved that if R is a strongly clean ring and $e^2 = e \in R$, then the corner ring eRe is strongly clean. For strongly $g(x)$-clean rings, we have the following result:

Theorem 3.6. *Let R be a strongly $(x - a)(x - b)$-clean ring with $a, b \in C(R)$. Then for any $e^2 = e \in R, eRe$ is strongly $(x - ea)(x - eb)$-clean. In particular, if $g(x) \in$*

$(x-ea)(x-eb)C(R)[x]$ and R is strongly $(x-a)(x-b)$-clean with $a,b \in C(R)$, then eRe is strongly $g(x)$ -clean.

Proof. By Theorem 2.4, R is strongly $(x-a)(x-b)$-clean if and only if R is strongly clean and $b-a \in U(R)$. If R is strongly clean, then eRe is strongly clean by [13]. Again by Theorem 2.4, eRe is strongly $(x-ea)(x-eb)$-clean. □

However, generally, strongly $g(x)$-clean property is not a Morita invariant: When $g(x) = (x-a)(x-b)$ where $a,b \in C(R)$ with $b-a \in U(R)$, the matrix ring over the local ring $\mathbb{Z}_{(p)}$ is not strongly clean [3] (hence, not strongly $g(x)$-clean).

4. STRONGLY $g(x)$-CLEAN RINGS VS RINGS GENERATED BY UNITS AND ROOTS OF 1

For any $n \in \mathbb{N}$, $U_n(R)$ denotes the set of elements of R which can be written as a sum of no more than n units of R [8]. A ring R is called *generated by its units* if $R = \bigcup_{n=1}^{\infty} U_n(R)$. We use strong $g(x)$-cleanness to characterize some rings in which every element can be written as the sum of unit and a root of 1 which commute.

Theorem 4.1. *Let R be a ring and $n \in \mathbb{N}$. Then the following are equivalent:*

1. *R is strongly $(x^2 - 2^n x)$-clean.*
2. *R is strongly (x^2-1)-clean.*
3. *R is strongly clean and $2 \in U(R)$.*
4. *$R = U_2(R)$ and for any $a \in R$, a can be expressed as $a = u+v$ with some $u, v \in U(R), uv = vu$, and $v^2 = 1$.*

Proof. (1) ⇒ (3). To prove $2 \in U(R)$. Suppose $2 \notin U(R)$, then $\overline{R} = R/(2^n R) \neq 0$. Let $2^n = s + u$ with $s^2 - 2^n s = 0, u \in U(R)$, and $su = us$. $\overline{0} = \overline{2^n} = \overline{s} + \overline{u}$ implies that $\overline{s} = -\overline{u} \in U(\overline{R})$. But $\overline{s}^2 = \overline{s^2} = \overline{2^n s} = \overline{0}$, a contradiction. So $2 \in U(R)$. Let $a = 0$ and $b = 2^n$. Then by (1) of Theorem 2.4, R is strongly clean.

(3) ⇒ (1). By (1) of Theorem 2.4, R is strongly $(x^2 - 2^n x)$-clean.

(3) ⇒ (4). Let $a \in R$. By "(1) ⇔ (3)", let $n = 1$. Then $1 - a = s + u$ where $s^2 = 2s, u \in U(R)$, and $su = us$. Then $a = (-u) + (1-s)$ with $-u \in U(R), (1-s)^2 = 1$, and $(-u)(1-s) = (1-s)(-u)$.

(4) $\Rightarrow$ (3). Let $a \in R$. By (4), $1 - a = u + v$ where $u \in U(R), v^2 = 1$, and $uv = vu$. Thus, $a = (-u) + (1 - v)$ with $-u \in U(R), (1 - v)^2 = 2(1 - v)$, and $(-u)(1 - v) = (1 - v)(-u)$. By "(1) $\Leftrightarrow$ (3)" and $n = 1$, we proved that (4) implies (3).

(2) $\Rightarrow$ (4). If R is strongly $(x^2 - 1)$-clean, then for any $r \in R$, there exist $v, u \in U(R)$ such that $r = v + u$ with $v^2 = 1$ and $uv = vu$.

(4) $\Rightarrow$ (2). Let $a \in R$. Then a can be expressed as $a = u + v$ with $u, v \in U(R), v^2 = 1$, and $uv = vu$. So v is the root of $x^2 - 1$. Hence, R is strongly $(x^2 - 1)$-clean. □

Example 4.2. *Rings in Example 2.7 are strongly $(x^2 - nx)$-clean. In particular, they are strongly $(x^2 - 2^n x)$-clean rings in which every element can be written as the sum of a unit and a square root of* 1 *which commute.*

Example 4.3. *Let F be a field and V be a vector space over F of infinite dimension, and let R be the subring of $E = End_F(V)$ generated by the identity and the finite rank transformations. Then R is strongly clean* [13, Example 7]. *In fact, E is locally Artinian. So the matrix ring $\mathbb{M}_k(E)$ is strongly $(x^2 - nx)$-clean with* $\mathrm{char} F \nmid n$. *If* $\mathrm{char} F \neq 2$, *then every element in the matrix ring can be written as the sum of a unit and a square root of* 1 *which commute.*

Example 4.4. *Let $A = F[x_1, x_2, \cdots]$ be the polynomial ring in a countably infinite set of indeterminates $(x_1, x_2, \cdots)$ over a field F, and let $I = (x_1^{k_1}, x_2^{k_2}, x_3^{k_3}, ...)$ with $k_i > 0$. Then $R = A/I$ is a local ring of dimension* 0 *which is not Noetherian. But R is locally Artinian. So the matrix ring $\mathbb{M}_k(R)$ is strongly $(x^2 - nx)$-clean with* $\mathrm{char} F \nmid n$. *If* $\mathrm{char} F \neq 2$, *then every element in the matrix ring can be written as the sum of a unit and a square root of* 1 *which commute.*

Proposition 4.5. *Let R be a ring with $c, d \in C(R)$ and $d \in U(R)$. If R is strongly $(x^2 + cx + d)$-clean, then $R = U_2(R)$. In particular, if R is strongly $(x^2 + x + 1)$-clean, then $R = U_2(R)$ is strongly $(x^4 - x)$-clean with every element is the sum of a unit and a cubic root of* 1 *which commute with each other.*

Proof. The first statement is clear. Let $r \in R$. Then $r = s + u$ with $u \in U(R)$, $s^2 + s + 1 = 0$, and $su = us$. So $s^4 - s = 0$. Thus, R is strongly $(x^4 - x)$-clean. Moreover, ever element in strongly (x^2+x+1)-clean ring R can be written as the sum of a unit and a cubic root of 1 which commute with each other. ☐

Lemma 4.6. [5] *Let $a \in R$. The following are equivalent for $n \in \mathbb{N}$:*

(1) $a = a(ua)^n$ *for some* $u \in U(R)$.
(2) $a = ve$ *for some* $e^{n+1} = e$ *and some* $v \in U(R)$.
(3) $a = fw$ *for some* $f^{n+1} = f$ *and some* $w \in U(R)$.

Proposition 4.7. *Let R be an strongly $(x^n - x)$-clean ring where $n \geq 2$ and $a \in R$. Then either (i) $a = u + v$ where $u \in U(R), v^{n-1} = 1$, and $uv = vu$ or (ii) both aR and Ra contain non-trivial idempotents.*

Proof. Since R is strongly $(x^n - x)$-clean, $a = s + u$ with $u \in U(R), s^n = s$, and $su = us$. Then $s^{n-1}a = s^{n-1}u + s$. So $(1 - s^{n-1})a = (1 - s^{n-1})u$. Since $1 - s^{n-1}$ is an idempotent, by Lemma 4.6, $(1 - s^{n-1})u = vg$ where $v \in U(R)$ and $g^2 = g \in R$. So $g = v^{-1}(1 - s^{n-1})a \in Ra$. Suppose (i) does not hold, then $1 - s^{n-1} \neq 0$, this implies $g \neq 0$. Thus, Ra contains a non-trivial idempotent. Similarly, aR contains a non-trivial idempotent. ☐

Finally, we give a property which has nothing to do with rings generated by units but it relates to strongly $(x^n - x)$-clean rings.

Proposition 4.8. *Let R be a ring and $n \in \mathbb{N}$. Then R is strongly $(ax^{2n} - bx)$-clean if and only if R is strongly $(ax^{2n} + bx)$-clean.*

Proof. " $\Rightarrow$ ". Suppose R is strongly $(ax^{2n} - bx)$-clean. Then for any $r \in R$, $-r = s + u$, $as^{2n} - bs = 0, u \in U(R)$, and $su = us$. So $r = (-s) + (-u)$ where $(-u) \in U(R), a(-s)^{2n} + b(-s) = 0$, and $(-s)(-u) = (-u)(-s)$. Hence, r is strongly $(ax^{2n} + bx)$-clean. Therefore, R is strongly $(ax^{2n} + bx)$-clean.

" $\Leftarrow$ ". Suppose R is strongly $(ax^{2n} + bx)$-clean. Let $r \in R$. Then there exist s and u such that $-r = s + u$, $as^{2n} + bs = 0, u \in U(R)$, and $su = us$. So $r = (-s) + (-u)$ satisfies $a(-s)^{2n} - b(-s) = 0, -u \in U(R)$, and $(-s)(-u) = (-u)(-s)$. Hence, R is strongly $(ax^{2n} - bx)$-clean. ☐

For $2n + 1 \in \mathbb{N}$, we do not know if the strong $(x^{2n+1} - x)$-cleanness of R is equivalent to the strong $(x^{2n+1} + x)$-cleanness of R.

Acknowledgments

The first author is partially supported by NSERC, Canada and the second is supported by the Research Grant from Harbin Institute of Technology.

References

[1] F. Azarpanah, When is $C(X)$ a clean ring? *Acta Math. Hungar.*, **94** (1-2) (2002): 53-58.

[2] V. P. Camillo and J. J. Simón, The Nicholson-Varadarajan theorem on clean linear transformations, *Glasgow Math. J.*, **44** (2002): 365-369.

[3] J. Chen, X. Yang, and Y. Zhou, On strongly clean matrix and trianglular matrix rings, *Comm. Algebra*, **34** (10) (2006): 3659-3674.

[4] W. Chen, A question on strongly clean rings, *Comm. Algebra*, **34** (7) (2006): 2374-2350.

[5] L. Fan and X. Yang, On rings whose elements are the sum of a unit and a root of a fixed polynomial, *Comm. Algebra*, **36** (1) 2008: 269-278.

[6] L. Fan and X. Yang, Strongly clean property and stable range one of some rings, preprint.

[7] J. Han and W. K. Nicholson, Extensions of clean rings, *Comm. Algebra*, **20** (2001): 2589-2596.

[8] M. Henriksen, Two classes of rings generated by their units, *J. Algebra*, **31** (1974): 182-193.

[9] W. W. McGovern, Clean semiprime f-rings with bounded inversion, *Comm. Algebra*, **31** (7) (2003): 3295-3304.

[10] W. K. Nicholson, Lifting idempotents and exchange rings, *Trans. Amer. Math. Soc.*, **229** (1977): 269-278.

[11] W. K. Nicholson, Strongly clean rings and Fitting's lemma, *Comm. Algebra*, **27** (1999): 3583-3592.

[12] W. K. Nicholson and Y. Zhou, Endomorphisms that are the sum of a unit and a root of a fixed polynomial, *Canad. Math. Bull.*, **49** (2006): 265-269.

[13] Sánchez Campos, On strongly clean rings, 2002, unpublished.

[14] Z. Wang and J. Chen, A note on clean rings, *Algebra Colloquium*, **14** (3) (2007): 537-540.